

Meta-Analysis

> Abdom Radiol (NY). 2025 Jan;50(1):198-212. doi: 10.1007/s00261-024-04302-y.

Epub 2024 Jul 26.

Qualitative and quantitative differentiation efficiency of dual-tracer PET/CT with 18F-fluorodeoxyglucose and ¹¹C-acetate for primary hepatocellular carcinoma: a systematic review and meta-analysis

Alisa Mohebbi ^{1 2}, Iman Kiani ^{1 2}, Saeed Mohammadzadeh ^{1 2}, Afshin Mohammadi ³,
Seyed Mohammad Tavangar ^{4 5}

Affiliations

— collapse

Affiliations

- ¹ Universal Scientific Education and Research Network (USERN), Tehran, Iran.
- ² School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.
- ³ Department of Radiology, Faculty of Medicine, Urmia University of Medical Science, Urmia, Iran.
- ⁴ Department of Pathology, Dr. Shariati Hospital, Tehran University of Medical Sciences, Tehran, Iran. tavangar@ams.ac.ir.
- ⁵ Chronic Diseases Research Center, Endocrinology and Metabolism Population Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran. tavangar@ams.ac.ir.

PMID: 39060514 DOI: [10.1007/s00261-024-04302-y](https://doi.org/10.1007/s00261-024-04302-y)

Abstract

FULL TEXT LINKS



ACTIONS

- “

Cite
- 🔖

Collections
- ↪

Permalink

PAGE NAVIGATION

< Title & authors

Abstract

Conflict of interest statement

Similar articles

Cited by

References

Publication types